



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2021

The first fossil record of the Sawshark *Pliotrema* (Pristiophoridae) from the Neogene of the South-East Pacific (Chile)

Guicharrousse-Vargas, Maurice ; Villafañá, Jaime ; Carrillo-Briceño, Jorge ; Oyanadel-Urbina, Pablo ; Figueroa, Romina ; Pérez-Marín, José ; Rivadeneira, Marcelo ; Kriwet, Jürgen

Abstract: We present the first fossil record of the sawshark genus *Pliotrema* from the south-eastern Pacific Ocean. The examined material was obtained from a little-known fossil locality named “Arenas de Caldera” in the Atacama region of Chile. The fossiliferous deposits belong to the Bahía Inglesa Formation, which is most likely middle Miocene–early Pliocene in age. There are no extant species of this sawfish in the eastern Pacific, probably due to the onset of cooling conditions during the Neogene. The type of environment for the elasmobranchs reported in this study is interpreted as demersal, based on the bathymetric affinities of extant species. Our results show that future studies are needed to better understand the evolutionary history and past distributions of this sawshark genus and their implications on current biogeographic patterns.

DOI: <https://doi.org/10.5710/amgh.01.03.2021.3389>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-203039>

Journal Article

Accepted Version

Originally published at:

Guicharrousse-Vargas, Maurice; Villafañá, Jaime; Carrillo-Briceño, Jorge; Oyanadel-Urbina, Pablo; Figueroa, Romina; Pérez-Marín, José; Rivadeneira, Marcelo; Kriwet, Jürgen (2021). The first fossil record of the Sawshark *Pliotrema* (Pristiophoridae) from the Neogene of the South-East Pacific (Chile). *Ameghiniana*, 58(2):122-131.

DOI: <https://doi.org/10.5710/amgh.01.03.2021.3389>



AMEGHINIANA

A GONDWANAN PALEONTOLOGICAL JOURNAL



This file is an uncorrected accepted manuscript (i.e., postprint). Please be aware that during the production process this version will definitively change. This postprint will be removed once the paper is officially published.

All legal disclaimers that apply to the journal pertain.

Submitted: August 20th, 2020 – **Accepted:** March 1st, 2021 – **Posted online:** March 8th, 2021

To link and cite this article:

doi: 10.5710/AMGH.01.03.2021.3389

PLEASE SCROLL DOWN FOR ARTICLE

THE FIRST FOSSIL RECORD OF THE SAWSHARK *PLIOTREMA*
(PRISTIOPHORIDAE) FROM THE NEOGENE OF THE SOUTH-EASTERN
PACIFIC (CHILE)

PRIMER REGISTRO FÓSIL DEL TIBURÓN SIERRA *PLIOTREMA*
(PRISTIOPHORIDAE) DEL NEÓGENO DEL PACÍFICO SUDESTE (CHILE)

MAURICE GUICHARROUSSE-VARGAS¹, JAIME A. VILLAFANA^{2,3}, JORGE D.
CARRILLO-BRICEÑO⁴, PABLO OYANADEL-URBINA⁵, ROMINA FIGUEROA¹, ,
JOSÉ R. PÉREZ-MARÍN⁶, MARCELO M. RIVADENEIRA^{5,7,8} AND JÜRGEN
KRIWET²

¹Museo Paleontológico de Caldera, Ilustre Municipalidad de Caldera, Cousiño 395,
1570000 Caldera, Chile. mguicharrousse@caldera.cl ; rfigueroa@caldera.cl

²University of Vienna, Department of Palaeontology, Althanstraße 14, Geocenter, 1090
Vienna, Austria. villafanaj88@univie.ac.at; juergen.kriwet@univie.ac.at

³Centro de Investigación en Recursos Naturales y Sustentabilidad, Universidad
Bernardo O'Higgins, Avda. Viel 1497, 8370993 Santiago, Chile.

⁴Paläontologisches Institut und Museum, Universität Zürich, Karl-Schmid-Straße 4,
8006 Zürich, Switzerland. jorge.carrillo@pim.uzh.ch

⁵Centro de Estudios Avanzados en Zonas Áridas (CEAZA), Av. Ossandón 877,
1781681 Coquimbo, Chile. pablo.oyanadel@ceaza.cl; marcelo.rivadeneira@ceaza.cl

⁶Oficina Técnica Regional del Consejo de Monumentos Nacionales en Magallanes,
O'Higgins 498, 6200000 Punta Arenas, Chile. jperez@monumentos.gob.cl

22 ⁷Departamento de Biología Marina, Facultad de Ciencias Biológicas, Universidad
23 Católica del Norte, Larrondo, 1281, 1781421 Coquimbo, Chile

24 ⁸Departamento de Biología, Universidad de La Serena, Raúl Bitrán 1305, 1720256 La
25 Serena, Chile.

26 22 pag.; 5 figs.; 1 table

27 GUICHARROUSE-VARGAS *ET AL.*: FOSSIL SAWSHARK FROM SOUTH
28 AMERICA

29 Short Description: Paleobiogeographic and paleoecological implications of a fossil
30 specimen of *Pliotrema* from the eastern Pacific of South America

31 Corresponding author: Jaime A. Villafaña, villafanaj88@univie.ac.at

32

33

34

35

36

37

38

39

40

Abstract. We present the first fossil record of the sawshark genus *Pliotrema* from the south-eastern Pacific Ocean. The examined material was obtained from a little-known fossil locality named "Arenas de Caldera" in the Atacama region of Chile. The fossiliferous deposits belong to the Bahía Inglesa Formation, which is most likely of middle Miocene-early Pliocene age. There are no extant species of this sawfish in the eastern Pacific, probably due to the onset of cooling conditions during the Neogene. The type of environment for the elasmobranchs reported in this study is interpreted as demersal, based on the bathymetric affinities of extant species. Our results show that future studies are needed to better understand the evolutionary history and past distributions of this sawshark genus and their implications on current biogeographic patterns.

Keywords. Bahía Inglesa. Formation. South America. Elasmobranchii. Pristiophoridae.

Resumen. PRIMER REGISTRO FÓSIL DEL TIBURON SIERRA *PLIOTREMA* (PRISTIOPHORIDAE) PARA EL NEÓGENO DEL PACÍFICO SUDESTE (CHILE).
Nosotros presentamos el primer registro fósil del tiburón sierra del género *Pliotrema* para el Pacífico sudeste. El material examinado fue obtenido de una localidad poco conocida llamada “Arenas de Caldera” en la región de Atacama en Chile. Los depósitos fosilíferos forman parte de la Formación Bahía Inglesa, la cual posee una edad probable de Mioceno medio a Plioceno temprano. No existen especies actuales de este tiburón sierra el Pacífico este, probablemente debido al establecimiento de las condiciones frías del Neógeno. El tipo de ambiente para los elasmobranchios reportados en este estudio es interpretado como demersal, basado en las afinidades batimétricas de las especies actuales. Nuestros resultados muestran que estudios futuros son necesarios para entender la historia evolutiva y distribución en el pasado de este tiburón sierra y las implicancias en sus patrones biogeográficos actuales.

Palabras clave. Formación Bahía Inglesa. América del sur. Chile. Elasmobranchii. Pristiophoridae.

SEVERAL CHONDRICHTHYAN TAXA have been reported from Neogene geological units along the Chilean coast up to now (*e.g.*, Suárez & Marquardt, 2003; Carrillo-Briceño *et al.*, 2013; Staig *et al.*, 2015; Villafaña & Rivadeneira, 2014, 2018; Partarrieu *et al.*, 2018). Along the eastern Pacific coast of Chile, the Bahía Inglesa Formation (Fm.) is one of the well-known Neogene units where chondrichthyan and bony fish remains are abundant, with fossiliferous localities ranging from the middle Miocene to the Pliocene (Walsh, 2001; Villafaña *et al.*, 2019). Despite the large number of chondrichthyan remains recovered from the Bahía Inglesa Fm., as well as from other Neogene units of Chile throughout the 20th century, remains of the sawshark genus *Pliotrema* Regan, 1906 have not yet been reported. Villafaña and Rivadeneira (2014) were the first to mention the presence of *Pliotrema* fossils from the south-eastern Pacific (from the Atacama Region), but without providing figures or descriptions regarding these remains. The worldwide fossil record of *Pliotrema* is also scarce (Cappetta, 2012), and includes a few reports from the late Paleocene of California (Welton, 1974; not illustrated) and middle Miocene of South Africa (Case, 1973). Here, for the first time, we illustrate, describe and discuss the geological, paleobiogeographic and paleoecological implications of the presence of a fossil specimen of *Pliotrema*. The occurrence of *Pliotrema* in the Neogene of Chile fills an exceptional gap in the distributional and evolutionary history of this sawshark genus.

Institutional abbreviation. MPC, Museo Paleontológico de Caldera, Caldera, Chile.

Anatomical abbreviations. Bh, barbed hook; Cap, enameloid part of a rostral spine;

LSce, less serrated cutting edge; Pdl, peduncle.

GEOLOGICAL SETTINGS

The Bahía Inglesa Formation (middle Miocene–lower Pleistocene) outcropping along the coastline of the Atacama region in Chile is characterized by a stratigraphic section

that correlates different units from the upper to lower coast, with oscillations in sea level combined with tectonics movements (Le Roux *et al.*, 2016). Lithologically, it is represented by a domain of coquinoid sandstones, marls and mudstones. The mudstones include phosphorites, diatomites and volcanic ash layers (Rojo, 1985; Godoy *et al.*, 2003). The fossiliferous locality “Arenas de Caldera”, from where the fossil described here comes (MPC-217), is exposed in the community of Caldera, which forms part of an extensive emerged coastal platform. It extends 20 km in a NE-SW direction and contains at least six relicts of coastal terraces that are distributed from sea level to 230 m.a.s.l (Guicharrousse *et al.*, 2018).

The Arenas de Caldera locality (Fig. 1; 27°4'37.62''S; 70°48'57.84''E) was mentioned in previous studies as being part of the Bahía Inglesa Fm. (Walsh & Suárez, 2005). This fossiliferous locality is mainly characterized by a sandstone sequence, where three well-defined stratigraphic horizons are distinguished (Fig. 2). Specimen MPC-217 was found in the second horizon, which is approximately 66 cm thick and represents a normal gradation that ranges from medium to very coarse sandstones with cross stratification.

Chronostratigraphy of the Bahía Inglesa Formation

According to Le Roux *et al.* (2016), the Bahía Inglesa Fm. ranges from the middle Miocene to Pleistocene. Although the Arenas de Caldera locality has not been unambiguously dated up to now, the presence of some fossil species can be used to infer a probable age range. In the eastern Pacific of South America, chondrichthyan taxa are commonly reported from Miocene to Pliocene sediments (Villafaña & Rivadeneira, 2014, 2018). For example, the extinct species †*Cosmopolitodus hastalis* is usually reported from Miocene sediments in Chile and Perú (Ehret *et al.*, 2012; Valenzuela-Toro *et al.*, 2016). Although the species *Carcharodon carcharias* is used as an indicator

of Pliocene age (de Muizon & Dvries, 1985), it also was reported as “probable” in late Miocene localities of Chile (Suárez & Marquardt, 2003; Walsh & Martill, 2006; Valenzuela-Toro *et al.*, 2013). The only identified marine invertebrate from Arenas de Caldera locality layer is *Crassostrea* sp., which is very abundant in sediments ranging from 15.3 to 4.2 Ma in the Bahía Inglesa Fm. (Le Roux *et al.*, 2016). Based on the stratigraphic position and biochron of some vertebrate and invertebrate taxa, this fossiliferous locality could be tentatively assigned to the middle Miocene to early Pliocene ages. Nevertheless, future radiometric dating could help to improve this inference.

MATERIAL AND METHODS

The single specimen MPC-217, which represents a rostral spine, is housed in the Museo Paleontológico de Caldera (MPC), Chile. The information about the current global distribution of *Pliotrema* species was obtained from Weigmann *et al.* (2020). Spine terminology follows Cappetta (2012), and the systematics for fossil and recent taxa are consistent with Weigmann *et al.* (2020) and Cappetta (2012).

According to Smith *et al.* (2015), the saw-teeth of sawfishes and sawsharks are not homologous to oral teeth, because they do not have the same origin (*i.e.*, derived from dermal denticles). These modified dermal denticles are recently named as rostral spines in the description of a new species of sawshark (†*Pristiophorus laevis*) from the Eocene of Antarctica (Engelbrecht *et al.*, 2017). We followed these recent studies and named our specimen MPC-217 as a rostral spine instead of rostral teeth.

157
158 **SYSTEMATIC PALEONTOLOGY**

159 Class CHONDRICHTHYES Huxley, 1880

160 Subclass ELASMOBRANCHII Bonaparte, 1838

161 Infraclass NEOSELACHII Compagno, 1977

162 Order PRISTIOPHORIFORMES Berg, 1958

163 Family PRISTIOPHORIDAE Bleeker, 1859

164 Genus *Pliotrema* Regan, 1906

165 **Type species.** *Pliotrema warreni* Regan 1906. Recent.

166 *Pliotrema* sp.

167 Figure 3. 1–3.6

168 **Material.** One rostral spine (MPC-217).

169 **Geographic distribution.** Arenas de Caldera locality.

170 **Stratigraphic distribution.** Middle Miocene–early Pliocene of Bahía Inglesa Fm.

171 **Description.** The rostral spine (MPC-217) is long, slender, and dorso-ventrally
172 flattened, reaching 42 mm in length and 16 mm in width (Fig. 3.1–3.4). The crown
173 (including the neck separating the crown from the peduncle) is slightly more than 4
174 times larger than the peduncle. The posterior cutting edge is weakly serrated with the
175 serrations restricted to the upper portion of the enameloid cap, but not reaching the apex
176 (Fig. 3.1–3.3). The anterior cutting edge is smooth and continuous and without any
177 serrations (Fig. 3.4). The anterior and posterior cutting edges are slightly tapered
178 laterally. The apex is acute and slightly bent towards the rear. The basal peduncle is
179 slender but broader than the crown jutting slightly out all around the crown in apical
180 view. It is subrectangularly shaped in its upper portion but with posteriorly elongated
181 base giving it an asymmetrical outline in dorsal and ventral views (Fig. 3.1–3.2). An

elliptically and deeply elongated foramen opens in the basal, concave face of the peduncle (Fig. 3.5).

Remarks. Rostral spines of pristiophoriforms are readily distinguishable from those of pristiform sawfishes in the development of a distinct peduncle, which is necessary to anchor the spine to the lateral rostrum wall, whereas they are inserted deeply into sockets in sawfishes lacking this peduncle (Wueringer *et al.*, 2009) The only other known elasmobranch clade possessing rostral spines are extinct sclerorhynchoid sawfishes, which, however, went extinct during the K/Pg boundary event (Kriwet & Benton, 2004).

Pristiophoriform sawsharks include a single family, Pristiophoridae, to which two extant genera (*Pliotrema* and *Pristiophorus*) and one extinct genus (\dagger *Ikamanius*) are assigned (Cappetta, 2012; Froese & Pauly, 2019). Although the extant and extinct members of the family Pristiophoridae (\dagger *Ikamanius*, *Pliotrema* and *Pristiophorus*) display similar morphologies in their rostral spines (*i.e.*, long and slender enameloid cap with a basal peduncle), they can be easily distinguished (Cappetta, 2012; Weigmann *et al.*, 2014). The rostral spines of extant *Pristiophorus* completely lack any serration along the anterior or posterior cutting edge (Ebert & Cailliet, 2011; Ebert & Wilms, 2013; Engelbrecht *et al.*, 2017; Villafañá *et al.*, 2019), whereas in \dagger *Ikamanius*, the anterior and posterior cutting edges are strongly serrated (Keyes, 1979; Cappetta, 2012). The presence of posterior serrations on the rostral spine's cutting edges is a character present in all known extant *Pliotrema* species; only varying in their degree and number of serrations (Weigmann *et al.*, 2020; Fig. 4). Although the specimen MPC-217 from the Bahía Inglesa Fm. displays the serrations along the posterior cutting edge as in rostral spines of the genus *Pliotrema* (*i.e.*, posterior cutting edge), its posterior cutting-edge serrations differ from those known from both fossil specimens (see Case, 1979;

Cappetta, 2012) and extant species (see Weigmann *et al.*, 2020) (Fig. 4). The comparison of fossil and extant specimens shows that the general spine morphology resembles that of *Pliotrema*, but that the serration of the posterior cutting edge is distinctly different. Following descriptions by Weigmann *et al.* (2020), two types of serrations are distinguished along the posterior cutting edge: type 1 consisting of a less serrated cutting edge and type 2 characterized by a serration developed into barbed hooks. Type 1 is similar to the reduced serration displayed by the fossil specimen reported here (Fig. 4.1), whereas type 2 is present in both fossil (Case, 1973; Cappetta, 2012) and extant specimens (Weigmann *et al.*, 2020) (Fig. 4.2–4.6). It nevertheless is mandatory that the fossil record of rostral pristiphoriform spines, especially those assigned to *Pliotrema* is revised to establish whether this kind of serration also occurs in other extinct members of this genus.

It is noteworthy that species level identification based on rostral teeth is not possible due to the high degree of morphological variation exhibited by *Pliotrema* species (Herman *et al.*, 1992; Weigmann *et al.*, 2020; Fig. 4) similar to what has been shown also for *Pristiophorus* (e.g., Engelbrecht *et al.*, 2017). Consequently, it only is possible to assign specimen MPC-217 to the genus *Pliotrema*, but not to any species. Nevertheless, the single specimen is of utmost importance since it represents the first record of a fossil rostral spine that can be referred unambiguously to *Pliotrema* and thus adds to our knowledge about the diversity of Neogene elasmobranchs from the eastern Paleo-Pacific.

DISCUSSION AND CONCLUSIONS

Paleoenvironmental conditions

230 In the Arenas de Caldera locality, the horizon with the highest abundance of fossil
231 remains (*i.e.*, H2) (Tab. 1) exhibits a cross-stratification as the main sedimentary
232 structure. Additionally, the inverse variation of the grain size, starting from medium to
233 high coarseness, allows us to infer that this change occurred gradually. The sedimentary
234 composition can be associated with off-shore coastal environments such as those
235 described for the Bahía Inglesa Fm. at Playa Chorillos locality, as well as the area close
236 to Los Dedos (Quezada *et al.*, 2007). The horizon “H2” concentrates the greatest variety
237 of disarticulated and abraded vertebrate remains, with a high abundance of
238 elasmobranch teeth (Tab. 1), and other vertebrates preliminarily identified as bony
239 fishes, mammals (cetaceans, pinnipeds and sirenians), sea birds and reptiles
240 (crocodylians). Invertebrates from the same layer include *Crassostrea* sp. and abundant
241 indeterminate bivalves. The elasmobranch taxa identified in this layer have extant
242 relatives with a maximum depth distribution of around 200 m (*e.g.*, *Brachaelurus*,
243 *Carcharias*) and 400 m (*e.g.*, *Myliobatis*), respectively (Tab. 1). However, other sharks
244 inhabiting continental and insular slopes (*e.g.*, *Carcharhinus*, *Odontaspis*,
245 *Pristiophorus*) occur at depths of up to 1000 m depth (Tab. 1). In the case of *Pliotrema*,
246 the three extant species are demersal sharks, but also show different depth preferences,
247 from shallow to deep waters (Weigmann *et al.*, 2020). *Pliotrema annae* Weigmann,
248 Gon, Leeney & Temple, 2020 is associated with shallow waters (20–35m), *Pliotrema*
249 *kajae* Weigmann, Gon, Leeney & Temple, 2020 with upper insular slopes and
250 submarine ridges (214–320m), whereas *Pliotrema warreni* Regan, 1906 is a demersal
251 species inhabiting the continental shelf and upper slope (10–915 m) (Weigmann *et al.*,
252 2020). Nevertheless, according to Weigmann *et al.* (2020), the depth preferences of the
253 two recently described species (*P. annae* and *P. kajae*) cannot be reliably inferred due to
254 them being defined based on low numbers of known specimens. For instance, *P. annae*

possibly also occurs in deeper waters during the day but enters shallow water during the night. Consequently, the mixture of depth distributions of the known elasmobranch taxa reported from Arenas de Caldera locality and the unclear, but seemingly varied depth preferences of the genus *Pliotrema* preclude a more elaborated interpretation. Thus, we can only infer that the genus *Pliotrema* most likely inhabited a demersal environment during the Neogene, based on the vertical distribution of extant representatives (Weigmann *et al.*, 2020). Although, we represent here the first attempt to describe the paleoenvironmental conditions of the Arenas de Caldera locality, more information about its depositional conditions and associated fauna would allow a more accurate conclusion.

Paleobiogeographic implications

Pliotrema is represented by three extant species: *P. warreni*, *P. kaja*, and *P. annae*, with distributions in the southwestern Indian and southern Atlantic oceans (Weigmann *et al.*, 2020) (see Fig. 5). The species *P. annae* is known from off the coast of Zanzibar, *P. kaja* from off the coast of Madagascar and the Mascarene Ridge (Weigmann *et al.*, 2020), while *P. warreni* is known from the coast of South Africa and the coast of southern Mozambique (Weigmann *et al.*, 2020).

The oldest record of *Pliotrema* comes from the Paleocene of North America, corresponding to isolated rostral spines (*Pliotrema* sp.) collected in the Lodo Formation, a unit deposited in an outer sublittoral to upper bathyal environment with temperate water conditions (Welton, 1974). Although there is no clear evidence, *Pliotrema* likely could have migrated from North America to the southern region during the Neogene (Fig. 5). At this time, the genus could have also migrated from the eastern Pacific through the Central America Sea Way and proto-Caribbean to the Atlantic Ocean, and

279 then into the Western Indian Ocean. This hypothesis could be supported by the presence
280 of *Pliotrema* in the late Miocene of South Africa (Case, 1973). Today, extant species of
281 *Pliotrema* are absent along both eastern Pacific and western Atlantic coasts of the
282 Americas (Weigmann *et al.*, 2020). The extirpation of *Pliotrema* from both regions is
283 likely correlated with the climatic and oceanographic events that occurred in the last 20
284 Ma, especially those processes associated with the closure of the Central America Sea
285 Way (CAS) and the rise of the Panamanian Isthmus (see Carrillo-Briceño *et al.*, 2018),
286 the opening of the Drake passage (Scher & Martin, 2006), and the onset of cold coastal
287 upwelling cells of the Humboldt Current system (Dekens *et al.*, 2007). According to
288 Carrillo-Briceño *et al.* (2018), sharks were the most affected chondrichthyan group in
289 Tropical America, with at least 24 genera affected by extirpation/extinction processes.
290 The extirpation of the sawsharks (Pristiophoridae) from the eastern Pacific at the end of
291 the Neogene is also intriguing (Carrillo-Briceño *et al.*, 2018). *Pristiophorus* had a wider
292 distribution in the eastern Pacific and western Atlantic oceans during Miocene and
293 Pliocene times (Carrillo-Briceño *et al.*, 2016, appendix 1; 2018, table S3). However, the
294 only extant species inhabiting the continent (*Pristiophorus schroederi*) has a reduced
295 geographical distribution in the Bahamian region (between Cuba, Florida, and the
296 Bahamas) over continental and insular slopes between 400 and 1000 m (Kiraly, *et al.*,
297 2003; Nevatte & Williamson, 2020). According to Villafañá and Rivadeneira (2018),
298 the regional extirpation of chondrichthyan genera from the eastern Pacific of South
299 America (including Pristiophoridae) was caused by changes in salinity and sea surface
300 temperature. The limited thermal tolerance to cold-water conditions was suggested as
301 the main reason for the demise of Neogene sharks along the southeastern Pacific
302 (Partarrieu *et al.*, 2018). Still, other factors such as habitat loss due to sea level changes

(e.g., Cione *et al.*, 2007) also could have played an important role in structuring the Neogene elasmobranch associations along the American Pacific coast.

ACKNOWLEDGMENTS

Financial support for this study was provided by a Becas-Chile Scholarship, Advanced Human Capital Program of the National Commission for Scientific and Technological Research (CONICYT, Chile) to JAV and FONDECYT-Chile grant # 1200843 to MMR. We deeply appreciate the comments and help provided by Gerard. R. Case, Thomas Reinecke, Jürgen Pollerspöck. This article was greatly improved by the comments and suggestion made by O. Aguilera and one anonymous reviewer. We also are grateful to Thodoris Argyriou and Ana Mercedes Balcarcel, for suggestions and comments on an earlier version of the manuscript.

REFERENCES

- Berg, L. S. (1958). System der Rezenten und Fossilen Fischartigen und Fische. Hochschulbücher für Biologie, Berlin, 310 pp.
- Bleeker, P. (1859). Enumeratio specierum piscium hucusque in Archipelago indico observatarum. *Acta Societatis scientiarum Indo-Neerlandae*, 6, 1–276.
- Bonaparte, C. L. (1838). Selachorum tabula analytica, Systema Ichthyologicum. *Memoires de la Societe Neuchateloise des Sciences Naturelles*, 2, 1–16.
- Cappetta, H. (2012). *Handbook of Paleoichthyology, Vol. 3E: Chondrichthyes. Mesozoic and Cenozoic Elasmobranchii: Teeth*. Verlag Dr. Friedrich Pfeil.

- 323 Carrillo-Briceño, J. D., Argyriou, T, Zapata, V, Kindlimann, R, Jaramillo, C. A. (2016).
 324 A new Early Miocene (Aquitania) Elasmobranchii assemblage from the Gaujira
 325 Peninsula, Colombia. *Ameghiniana*, 53(2),77–99.
- 326 Carrillo-Briceño, J. D., Carrillo, J. D., Aguilera, O. A., & Sánchez-Villagra, M. R.
 327 (2018). Shark and ray diversity in the Tropical America (Neotropics)—an
 328 examination of environmental and historical factors affecting diversity. *PeerJ*, 6,
 329 e5313.
- 330 Carrillo-Briceño, J. D, González-Barba G., Landaeta M. F., & Nielsen SN. (2013).
 331 Condrictios fósiles del Plioceno Superior de la Formación Horcón, Región de
 332 Valparaíso, Chile central. *Revista Chilena de Historia Natural*, 86(2), 191–206.
- 333 Case, G. R. (1973). *Fossil sharks: a pictorial review*. Pioneer Litho Company.
- 334 Case, G. R. (1979). Collecting fossil shark's teeth at Big Brook, Monmouth County,
 335 New Jersey. *Bulletin of the Bergen County Mineral and Paleontological Society*,
 336 13(5), 12–14.
- 337 Cione, A.L., Mennucci, J., Santalucita, F., & Acosta Hospitaleche, C. (2007). Local
 338 extinction of sharks of genus *Carcharias* Rafinesque, 1810 (Elasmobranchii,
 339 Odontaspidae) in the eastern Pacific Ocean. *Revista Geológica de Chile*, 34(1),
 340 139–145.
- 341 Compagno, L. J. V. (1977). Phyletic relationships of living sharks and rays. *American*
 342 *Zoologist*, 17(2), 303–322.

- 343 de Muizon, C., & Devries, T. J. (1985). Geology and paleontology of late Cenozoic
344 marine deposits in the Sacaco area (Peru). *Geologische Rundschau*, 74(3), 547–
345 563.
- 346 Dekens, P.S., Ravelo, A.C., & McCarthy, M. D. (2007). Warm upwelling regions in the
347 Pliocene warm period. *Paleoceanography*, 22, PA3211
- 348 Ebert, D. A., & Cailliet, G.M. (2011). *Pristiophorus nancyae*, a new species of
349 Sawshark (Chondrichthyes: Pristiophoridae) from Southern Africa. *Bulletin of*
350 *Marine Science*, 87(3), 501–512.
- 351 Ebert, D. A., & Wilms, H. A. (2013). *Pristiophorus lanae* sp. nov., a new sawshark
352 species from the Western North Pacific, with comments on the genus
353 *Pristiophorus* Müller & Henle, 1837 (Chondrichthyes: Pristiophoridae). *Zootaxa*,
354 3752, 86–100.
- 355 Ehret, D. J., Macfadden, B. J., Jones, D. S., Devries, T. J., Foster, D. A., & Salas-
356 Gismondi, R. (2012). Origin of the white shark *Carcharodon* (Lamniformes:
357 Lamnidae) based on recalibration of the Upper Neogene Pisco Formation of Peru.
358 *Palaeontology*, 55, 1139–1153.
- 359 Engelbrecht, A., Mörs, T., Reguero, M. A., & Kriwet, J. (2017). A new sawshark,
360 *Pristiophorus laevis*, from the Eocene of Antarctica with comments on
361 *Pristiophorus lanceolatus*. *Historical Biology*, 29(6), 841–853.
- 362 Froese, R. & Pauly, D. (2019). Fish Stocks. Encyclopedia of Life Science. Elsevier.

- 363 Godoy, E., Marquardt, C., & Blanco, N. (2003). *Carta Caldera, Región Atacama.*
 364 *Carta geológica de Chile, Serie Geología Básica, escala 1:100.000.* Servicio
 365 Nacional de Geología y Minería, Chile.
- 366 Guicharrousse, M., Pérez, J. R., & Oyanadel-Urbina, P. (2018). Estudio y
 367 contextualización estratigráfica de los fósiles de las Arenas de Caldera, Fm. Bahía
 368 inglesa, Región de Atacama, Chile: Parte I Estratigrafía. *Actas de 1^º Congreso de*
 369 *Paleontológico de Chile* (pp. 101–103). Punta Arenas.
- 370 Herman, J., Hovestadt-Euler, M., & Hovestadt, D. C. (1992). Contributions to the study
 371 of the comparative morphology of teeth and other relevant ichthyodorulites in
 372 living superspecific taxa of Chondrichthyan fishes. Part A: Selachii. No. 4: Order:
 373 Orectolobiformes Families: Brachaeluridae, Giglymostomatidae, Hemiscylliidae,
 374 Orectolobidae, Parascylliidae, Rhiniodontidae, Stegostomatidae. Order:
 375 Pristiophoriformes - Family: Pristiophoridae Order: Squatiniformes - Family:
 376 Squatinidae. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique,*
 377 *Biologie*, 62, 193–254.
- 378 Huxley, T. H. (1880). On the application of the laws of evolution to the arrangement of
 379 the Vertebrata, and more particularly of the Mammalia. *Proceedings of the*
 380 *Zoological Society of London*, 1880, 649–662.
- 381 Keyes, I. W. (1979). Ikamauius, a new genus of fossil sawshark (Order Selachii: Family
 382 Pristiophoridae) from the Cenozoic of New Zealand. *New Zealand Journal of*
 383 *Geology and Geophysics*, 22(1), 125–129.
- 384 Kiraly SJ, Moore JA, Jasinski PH. 2003. Deepwater and other sharks of the US Atlantic
 385 ocean exclusive economic zone. *Marine Fisheries Review*, 65,1–20.

- 386 Kriwet, J., & Benton, M. J. (2004). Neoselachian (chondrichthyes, elasmobranchii)
 387 diversity across the cretaceous–tertiary boundary. *Palaeogeography,*
 388 *Palaeoclimatology, Palaeoecology*, 214(3), 181–194.
- 389 Le Roux, J. P., Achurra, L., Henríquez, Á., Carreño, C., Rivera, H., Suárez, M. E., &
 390 Gutstein, C. S. (2016). Oroclinal bending of the Juan Fernández Ridge suggested
 391 by geohistory analysis of the Bahía Inglesa Formation, north-central Chile.
 392 *Sedimentary Geology*, 333, 32–49.
- 393 Nevatte, R. J., & Williamson, J. E. (2020). The sawshark redemption: current
 394 knowledge and future directions for sawsharks (Pristiophoridae). *Fish and*
 395 *Fisheries*, 21(6), 1213–1237.
- 396 Partarrieu, D., VillafañA, J. A., Pinto, L., Mourgues, F. A., Oyanadel-Urbina, P. A.,
 397 Rivadeneira, M. M., & Carrillo-Briceño, J. D. (2018). Neogene ‘horn sharks’
 398 Heterodontus (Chondrichthyes: Elasmobranchii) from the Southeastern Pacific
 399 and their paleoenvironmental significance. *Ameghiniana*, 55(6), 651–667.
- 400 Quezada, J., Gonzalez, G., Dunai, T., Jensen, A., & Juez-Larre, J. (2007). Alzamiento
 401 litoral Pleistoceno del norte de Chile: edades 21Ne de la terraza costera más alta
 402 del área de Caldera-Bahía Inglesa. *Andean Geology*, 34(1), 81–96.
- 403 Regan, C. T. (1906). A classification of the selachian fishes. *Proceedings of the*
 404 *Zoological Society of London*, 722–758.
- 405 Rojo, M. (1985). Un aporte al conocimiento del Terciario marino: Formación Bahía
 406 Inglesa. *Actas de IV Congreso Geológico Chileno* (pp. 514–532). Antofagasta.

407 Scher, H. D., & Martin, E. E. (2006). Timing and climatic consequences of the opening
 408 of Drake Passage. *Science*, 312(5772), 428–430.

409 Smith, M. M., Riley, A., Fraser, G. J., Underwood, C., Welten, M., Kriwet, J., Pfaff, C.,
 410 & Johanson, Z. (2015). Early development of rostrum saw-teeth in a fossil ray
 411 tests classical theories of the evolution of vertebrate dentitions. *Proceedings of the*
 412 *Royal Society B: Biological Sciences*, 282(1816), 20151628.

413 Staig, F., Hernández, S., López, P., Villafaña, J. A., Varas, C., Soto, L. P., & Carrillo
 414 Briceño, J. D. (2015). Late Neogene elasmobranch fauna from the Coquimbo
 415 formation, Chile. *Revista Brasileira de Paleontologia*, 18(2), 261–272.

416 Suárez, M. E., & Marquardt, C. (2003). Revisión preliminar de las faunas de peces
 417 Elasmobranquios del Mesozoico y Cenozoico de Chile: su valor como indicadores
 418 cronoestratigráficos. *Actas de 10° Congreso Geológico Chileno* (pp.1–9).
 419 Concepción.

420 Valenzuela-Toro, A. M., Gutstein, C. S., Varas-Malca, R. M., Suárez, M. E., &
 421 Pyenson, N. D. (2013). Pinniped turnover in the south Pacific Ocean: new
 422 evidence from the Plio-Pleistocene of the Atacama Desert, Chile. *Journal of*
 423 *Vertebrate Paleontology*, 33(1), 1–8.

424 Valenzuela-Toro, A. M., Pyenson N. D., Gutstein C. S., & Suárez M. E. (2016). A new
 425 dwarf seal from the late Neogene of South America and the evolution of
 426 pinnipeds in the southern hemisphere. *Palaeontology*, 2(1), 101–115.

427 Villafañá J.A., Marramà G., Hernandez S., Carrillo-Briceño J. D., & Hovestadt, D.
 428 (2019). The Neogene fossil record of Aetomylaeus (Elasmobranchii,
 429 Myliobatidae) from the southeastern Pacific. *Journal of Vertebrate Paleontology*,
 430 39(1), e1577251.

431 Villafañá, J. A., & Rivadeneira, M. M. (2014). Rise and fall in diversity of Neogene
 432 marine vertebrates on the temperate Pacific coast of South America.
 433 *Paleobiology*, 40(4), 659–674.

434 Villafañá, J. A., & Rivadeneira, M. M. (2018). The modulating role of traits on the
 435 biogeographic dynamics of chondrichthyans from the Neogene to the present.
 436 *Paleobiology*, 44(2), 251–262.

437 Walsh, S. A. (2001). *The Bahía Inglesa Formation Bonebed: genesis and palaeontology*
 438 *of a Neogene Konzentrat Lagerstätte from north–central Chile* (PhD thesis,
 439 University of Portsmouth).

440 Walsh, S. A., & Martill, D. M. (2006). A possible earthquake-triggered mega-boulder
 441 slide in a Chilean Mio-Pliocene marine sequence: evidence for rapid uplift and
 442 bonebed genesis. *Journal of the Geological Society*, 163(4), 697–705.

443 Walsh, S. A., & Suárez, M. E. (2005). First post – Mesozoic record of Crocodyliformes
 444 from Chile. *Acta Palaeontologica Polonica*, 50(3), 595–600.

445 Weigmann, S. (2016). Annotated checklist of the living sharks, batoids and chimaeras
 446 (Chondrichthyes) of the world, with a focus on biogeographical diversity. *Journal*
 447 *of Fish Biology*, 88(3), 837–1037.

448 Weigmann, S., Stehmann, M. F.W., & Thiel, R. (2014). Contribution to the taxonomy
 449 and distribution of *Pristiophorus nancyae* (Elasmobranchii: Pristiophoriformes)
 450 from the deep western Indian Ocean. *Marine Biodiversity*, 44(2), 189–202.

451 Weigmann, S., Gon, O., Leeney, R. H., Barrowclift, E., Berggren, P., Jiddawi, N., &
 452 Temple, A. J. (2020). Revision of the sixgill sawsharks, genus *Pliotrema*
 453 (Chondrichthyes, Pristiophoriformes), with descriptions of two new species and a
 454 redescription of *P. warreni* Regan. *PLoS ONE*, 15(3), e0228791.

455 Welton, B. J. (1974). Preliminary note on the Paleocene elasmobranchs of the Lodo
 456 Formation, Fresno County, California. In Payne, M. B. (ed.). *The Paleogene of the*
 457 *Panoche Creek-Cantua Creek Area, central California* (pp. 91–91). SEPM.

458 Wueringer, B. E., Squire, L., & Collin, S. P. (2009). The biology of extinct and extant
 459 sawfish (Batoidea: Sclerorhynchidae and Pristidae). *Reviews in Fish Biology and*
 460 *Fisheries*, 19(4), 445–464.

461
 462
 463

FIGURE AND TABLE CAPTIONS

Figure 1. Location map of the Arenas de Caldera locality.

Figure 2. Stratigraphic section of the Arenas de Caldera locality.

Figure 3. Rostral spine of *Pliotrema* sp. (MPC-127) from Arenas de Caldera locality.


Views: **1**, ventral; **2**, dorsal; **3**, posterior; **4**, anterior; **5**, basal; and **6**, occlusal. Scale bar equals 1 cm.


Figure 4. Line drawings of fossil and extant *Pliotrema* rostral teeth in dorsal views. **1**, this study (MPC-127); **2**, *Pliotrema* sp. from Miocene of South Africa (Cappetta, 2012); **3**, *Pliotrema* sp. from late Miocene of South Africa (Case, 1973); **4**, extant *Pliotrema annae* from Papa Unguja, Tanzania (Weigman *et al.*, 2020); **5**, extant *Pliotrema warreni* from Eastern Cape, South Africa (Weigman *et al.*, 2020); **6**, extant *Pliotrema kajae* from southwestern Madagascar (Weigman *et al.*, 2020). The specimens 2 and 3 were originally illustrated without scale. Bh, barbed hook; Cap, enameloid part of a rostral spine; LSce, less serrated cutting edge; Pdl, peduncle. Scale bar equals 5 mm.

Figure 5. Map showing the geographic distribution of extant and fossil *Pliotrema*.



Fossil distribution based on Case (1973), Welton (1974) and Cappetta (2012). Extant distribution based on Weigmann *et al.* (2020).

484 **Table 1.** Type of environment and depth distribution of the elasmobranch taxa reported
485 from Arenas de Caldera locality. Information taken from Fishbase (Froese & Pauly,
486 2019), Nevatte and Williamson (2020) and Weigmann *et al.* (2020).












Table 1. Type of environment and depth distribution of the elasmobranch taxa reported from Arenas de Caldera locality. Information taken from Fishbase (Froese & Pauly, 2019), Nevatte and Williamson (2020) and Weigmann *et al.* (2020)

Order	Family	Fossil taxa	Environment	min depth (m)	max depth (m)
Orectolobiformes	Brachaeluridae	<i>Brachaelurus</i> sp.	D-Ra	4	217
Carcharhiniformes	Carcharhinidae	<i>Carcharhinus</i> sp.	D-Pn-Ra	0	1082
Lamniformes	Carcharhinidae	<i>Carcharias</i> sp.	D-Ra	1	232
Lamniformes	Lamnidae	<i>Carcharodon carcharias</i>	Po	0	1200
Pristiophoriformes	Pristiophoridae	<i>Pliotrema</i> sp.	D	10	430
Lamniformes	Lamnidae	† <i>Cosmopolitodus hastalis</i>	-	-	-
Lamniformes	Lamnidae	<i>Isurus</i> sp.	Po	0	1752
Myliobatiformes	Myliobatidae	<i>Myliobatis</i> sp.	Bp-D-Ra	0	422
Hexanchiformes	Hexanchidae	<i>Notorynchus</i> sp.	B	0	570
Lamniformes	Odontaspidae	<i>Odontaspis</i> sp.	Bp-Po	10	1015
Pristiophoriformes	Pristiophoridae	<i>Pristiophorus</i> sp.	D-Po-Pn	0	1240

Bp, benthopelagic; D, demersal; Ra, reef-associated; Pn, pelagic-neritic; Po, pelagic-oceanic.